

A randomised prospective and retrospective study of biological plating for tibial fractures

Salim A. Lad¹, P.V Naveen Kumar², Pradip B. Patil³, Shailesh Deshpande⁴, Adarsh Kumbar⁵, Ravindra Hattarki⁶

¹ Professor & Head, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

² Junior Resident, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

³ Associate Professor, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

⁴ Assistant Professor, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

⁵ Junior Resident, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

⁶ Assistant Professor, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

ABSTRACT

A series of 30 patients of proximal and distal tibial fractures treated with minimally invasive techniques have been reviewed after surgery to study the results of fractures of proximal and distal tibia with biological plating. We have scored proximal tibial fractures according to Rasmussen scale and distal tibial fractures by Tenny and Wiss scale. Out of 30 patients, 4 had excellent, 15 had good, 7 patient had fair, and 3 had poor results. Two patients had superficial wound infection and two patients had deep infections. This minimally invasive technique for treatment of distal tibial fractures proves to be a feasible and worthwhile method of stabilization while avoiding the severe complications associated with the more standard methods of internal or external fixation of these fractures.

Key words: Biological plating of tibia, MIPPO.

Introduction

Fractures of the tibia are notoriously difficult to treat, and traditional methods of fixation are often fraught with complications due to limited soft tissue, subcutaneous location and poor vascularity. Young males are more commonly involved with an average of 37 yrs. Open reduction and stable internal fixation allows axial alignment of the limb, permits early mobilisation and results in bony union from endosteal bone healing. However in the tibial fractures the surgical dissection required to achieve anatomical reduction evacuates the osteogenic fracture haematoma

and causes soft tissue stripping that may result in infection, wound necrosis and delayed or non union. A balance between anatomical reduction and soft tissue devitalisation is therefore required. Minimally invasive surgery gives that direct anatomical reduction with rigid fixation and maintenance of alignment by bridging the fracture without compression and result in secondary bone healing. Biological plating preserves the soft tissue envelope and the periosteum maintains arterial vascularity and therefore minimises the surgical trauma to the zone of injury. Stripping of the periosteum is thus avoided, the fragments remain integrated into the soft tissue, and healing occurs spontaneously by way of callus formation

The advantages of biological fixation are:

1. Soft tissue remains intact.
2. There is some mobility in the fracture zone which accelerates the callus formation.

*Correspondence

Dr. Salim A Lad

Professor & Head, Department of Orthopaedics, D.Y. Patil Medical College & Hospital, Kolhapur, Maharashtra, India

3. Reduced surgical time and tourniquet time along with smaller incision.

Methodology

In this prospective study, 30 patients of proximal distal Tibia fractures admitted in our institute were studied during the period of August 2013 to July 2015. Patients with closed fractures of proximal and distal Tibia were included in this study and fractures in children and open fractures and pts with fractures accompanied with neurovascular injury are excluded from this study.

These all fractures were classified according to AO Classification. We used LCP for both proximal and distal tibia fracture fixation. Mippo technique was used. For distal tibia if there was fibula fracture then it was fixed first to achieve length and then distal tibia fracture was fixed.

Post-Operative Care

Intravenous antibiotic therapy was used for 2-3 days postoperatively according to the wound condition.

Passive ankle and knee joint motion was allowed as soon as possible.

After the soft tissues healed and postoperative swelling was diminished, the patient was allowed non-weight-bearing ambulation with crutches.

Progressive weight-bearing was encouraged once there was radiographic evidence of callus formation.

Fracture healing was assessed in the outpatient clinic at 6 weeks and 3 months postoperatively.

Clinical evidence of septic complications, incision breakdown, and skin necrosis was recorded.

Follow-up, including clinical examination and radiographic examination, was performed once a month until the fractures achieved clinical union based on the standard criteria on (pain-free full weight-bearing).

Final evaluation was done for distal tibial fractures as per Teeny and Wiss clinical assessment criteria which is based on 100 points system and radiological assessment by ACFAS ankle score.

Final evaluation was done for proximal tibia fractures as per Rasmussen criteria which is based on 100 points.

RESULTS



Fig. 1 : Preop and post op xray of distal tibia fractures treated with biological plating



Fig 2 : Range of movement at ankle joint after distal tibia fracture treated with biological plating




Fig 3 : Preop and post op xrays of Proximal tibial fracture treated with biological plating



Fig 4 : Incision scar of biological plating done for proximal tibia fracture with full extension of knee And full flexion in cross legged position

Table 1 : Functional grading of the patients as per scale)

Sr. No	Grading	No of Cases	Percentage
1	Excellent	5	16.667
2	Good	15	50
3	Fair	7	23.33
4	Poor	3	10

Table 2: Type of union

Sr. No	Type of fracture	No	Union	Delayed Union	Non Union
1	Type 4 1 A	3	2	1	0
2	Type 41 B	3	2	1	0
3	Type 41 C	4	3	1	0
4	Type 43 A	18	14	3	1
5	Type 43 B	0	0	0	0
6	Type 43 C	2	1	0	1

Discussion

The results of operative treatment are dependent on the severity of the initial injury and the quality and stability of the reduction. The mechanism of injury, the status of the soft tissues and the degree of comminution affect the long term clinical result. The interval for radiological union was 12 to 20 weeks in our study in accordance to previous study. However, the most important factor is to achieve stable fixation and to allow early range of motion without unnecessary osseous and soft tissue devascularization. Minimally invasive techniques are based on principles of limited exposure, indirect reduction methods and limited contact between bone and implant. As a result of these principles this technique, as seen in present study, avoided major soft tissue complications and shortened the length of the patient's stay in the hospital. The bone healing was excellent with this type of fixation because the stresses were distributed over a longer segment of bone. This technique can be used in fractures where locked nailing cannot be done like vertical slit and markedly comminuted fractures. There was rapid fracture consolidation due to preserved vascularity. There were fewer incidences of delayed union and non-union. There was decreased need for bone grafting. There was less incidence of infection due to limited exposure. There were less chances of refracture. There was no chance of vascular complication by carefully inserting the plate submuscularly through limited incisions. There was no need of any specialized instrumentation and the method was less time consuming and cost effective in the present study in a similar fashion as reported by previous authors. With the introduction of Locking Compression Plates (LCP), minimally invasive techniques have become widely used. The plates act as internal fixators in a bridging manner, thus resulting in secondary bone healing. Thus, Biological plating for tibial fractures will prove

to be a feasible and worthwhile method of stabilization while avoiding the severe complications associated with the more standard methods of internal or external fixation.

Conclusion

In this study we looked at the results and complications in 30 consecutive patients treated with percutaneous plating for fractures of the distal and proximal tibia with a minimum follow-up period of one year. No significant soft tissue problems occurred. The need for bone grafting should be carefully evaluated in every case as we encountered 6 delayed unions. All fractures healed within one year except 2 which went into non-union; there were 2 malunions. The use of indirect reduction techniques and small incisions to insert hardware is technically more demanding and requires strict radiographic control throughout the procedure, but it considerably decreases surgical trauma to the soft tissues. Most of the patients were in age group of 20-40 years (70%) with mean age of 36 years. Road traffic accidents were found to be the commonest mode of trauma (50%). Right limb was involved more often (70%) than the left. The mean time taken for starting partial weight bearing was 6 weeks, time for starting full weight bearing 12 to 22 weeks. The time taken for radiological union was 12 to 24 weeks. The average range of motion at the ankle for distal tibial fractures dorsi & plantar flexion were 5 to 15⁰ and 15to30⁰ respectively. The average range of motion at knee for proximal tibial fractures were flexion 60 to 135⁰ degrees and full extension and in 4 patients we got extension lag. According to Tenny & Wiss and Ramuseen criteria we got 4 excellent, 15 good, 7 fair and 3 poor results. In our series 2 patients

got superficial infection, 2 patients got deep infections and one patient had implant failure.

Acknowledgement

It gives us great pleasure to express our deepest sense of gratitude and thanks to the Dept. of Orthopaedics, D. Y. Patil Medical College, Kolhapur for helping us during this study.

References

1. Minimal fracture fixation by Bernhard g weber 2004 pages 24 – 36
2. AO principles of fracture management by Thomas P Ruedi William m Murphy 2000 page 221-230.
3. Textbook of Orthopaedics And Trauma by Dr G.S.Kulkarni 2nd Edition 2008 .2 (3):1441-1445 ;2138-2148
4. Whiteside LA, Lesker PA. The effects of extra- periosteal and subperiosteal dissection on fracture healing. *J Bone Joint Surg Am* 1978 ; 60 : 26-30.
5. Textbook of Orthopaedics And Trauma by Dr G.S.Kulkarni 2nd Edition 2008 Vol-3 2138-2148.
6. Peren SM.the concept of biological plating using the (LC-DCP) Scientific Background design and application.*Injury* 1991;1(1):1-41
7. Macnab I. The role of periosteal blood supply in healing of fracture of tibia. *Clinical Ortho*1974;105:27
8. Blauth M, Bastian L, Krettek C, Knop C, Evans S.Surgical options for the treatment of severe tibial pilon fractures: a study of three techniques. *J Orthop Trauma* 2001 ; 15 : 153-160.
9. Collinge CA, Sanders R. Percutaneous plating in the lower extremity. *J Am Acad Orthop Surg* 2000 ; 8 : 211-216
10. Leach IRE: A means of stabilizing comminuted distal tibial fractures. *J Trauma* 4: 722–730, 1964.
11. Rouff AC III, Snider RK: Explosion fractures of the distal tibia with major articular involvement. *J Trauma* 11: 866–871, 1971
12. Reudi T, Allgower M: The operative treatment of intraarticular fractures of the lower end of the tibia. *Clin Orthop*.1979; 138: 105–110
13. McFerran M, Smith S, Boulas HJ, Schartz H: Complications encountered in the treatment of pilon fractures. *J Orthop Trauma*.1992; 6: 195–200
14. Hazarikaa S Chakravarthy J , Cooper J . Minimally invasive locking plate osteosynthesis for fractures of distal tibia .*Injury* 2006; 37: 877-87.
15. Hasenboehler E , Rikli E , Babst R . Locking compression plate with minimally invasive plate osteosynthesis in diaphyseal and distal tibial fracture : A retrospective study of 32 patients . *Injury* 2007 ; 38: 365-70

Source of Support: Nil

Conflict of Interest: None