
A study on Variations of pulmonary function tests in different stages of pregnancy

N Bhaskar Rao*, Ch Rambabu

Department of Obstetrics & Gynaecology, Great Eastern Medical School and Hospital, Srikakulam, Andhra Pradesh, India

ABSTRACT

Pulmonary status of women during pregnancy is of great importance. Their efficacy should be assessed for both maternal & fetal well being. In this study, the respiratory functions in various stages of pregnancy along with postnatal status were monitored & compared. Mini Wright Peak Flow Meter (PFM), which is highly effective in assessing the lung functions of a person, has been selected as a means of assessing pulmonary functions of a pregnant woman. Graphs or tables are available of predicted normal values based on person's gender, age & height. We have tested the PEFR throughout pregnancy from 6 weeks onwards till puerperium and effect of mode of delivery and puerperium on it. We have taken study sample size of 300. It is divided as 60 subjects in 5 groups like Non pregnant, First trimester, Second trimester, Third trimester, Postnatal period till 30 days after delivery in medical teaching institute. Predicted PEFR which depends on age & height of patient is calculated by referring to Wright's scale. In 1st & 2nd trimester the lung functions are relatively normal, while in 3rd trimester the pulmonary function gets compromised with decrease in PEFR, PEFR is not affected by parity and in all modes of delivery it is significantly low, more after caesarean section (post operative pain might be responsible) and by the end of one month after delivery, the lung functions return back to normal.

Key words: Bupivacaine, Supraclavicular brachial plexus block, Peripheral nerve blockade.

Introduction

Giving birth to a child is a very precious event in the life of every woman. It is a time of celebrations in the family to welcome a new member. The pregnant woman goes through various physical, mental, psychological & social changes. The physiological changes occurring in a pregnant woman are vast & widespread. These include changes in genital organs, breasts, weight gain, Cutaneous changes, systemic changes (respiratory, cardiovascular, etc.), body water metabolism, hematological & metabolic changes. In first trimester, there is a feeling of nausea and vomiting while in later months, women mostly

complain of dyspnoea and palpitations due to mechanical discomfort caused by enlarged gravid uterus. We used Mini Wright Peak Flow Meter to assess the peak flow rate. It measures how fast a person can breathe out (exhale) air i.e. Peak Expiratory Flow Rate (PEFR). It provides an indirect assessment of airway hyper responsiveness which correlates closely with degree of airway inflammation. It is also used for monitoring of severity of asthma. PEFR varies with age, height & sex of patient. Depending on these parameters, Predicted PEFR is calculated. Observed PEFR is obtained with peak flow meter. Normally, observed PEFR should be >80% predicted PEFR. If the PEFR is <80% it indicates pulmonary functions are compromised. Likewise, the respiratory functions in various stages of pregnancy as well as in postnatal period are monitored and compared. The current study is to determine the pulmonary function test (PEFR) in different stages of antenatal (1st, 2nd, 3rd

*Correspondence

Dr. N Bhaskar Rao

Department of Obstetrics and Gynecology, Great Eastern Medical School and Hospital, Srikakulam, Andhra Pradesh, India

trimester) & postnatal period up to 6 weeks and to compare the pulmonary functions of non-pregnant & pregnant women.

Mini Wright peak flow meter

A peak flow meter is a medical device which measures and scales the peak expiratory flow rate (PEFR). PEFR is the maximum rate of airflow which can be achieved during a sudden forced expiration from a position of full inspiration. The Peak Flow Meter provides a

convenient, accurate & inexpensive method for monitoring lung function changes.

Indications: Estimation of airway caliber. Diagnosis, treatment, monitoring severity of asthma. To know asymptomatic airway obstruction, Evaluation of dyspnoea and peak flow monitoring at home. Clinical & epidemiological research studies. Establishment of baseline personal best.

Results

Table 1: Comparison of % of subjects in non-pregnant, pregnant & postnatal period

Subject	No. of Cases examined (300)	No subjects with PEFR Significantly low (93)	PEFR range & Mean
Non- pregnant	60	6(10.00%)	268-396(332)
First trimester	60	14(23.33%)	256-366(311)
Second trimester	60	28(46.66%)	224-308 (266)
Third trimester	60	36(60.00%)	142-266 (205)
Postnatal (immediate)	60	9(15.00%)	266-406 (336)

p<0.001, highly significant

Table 2: Effect of advancing gestation on % of subjects with significantly lowered PEFR

Gestational age (in weeks)	(78) No of subjects with PEFR significantly less
6-10	3%
11-15	6%
16-20	6%
21-25	8%
26-30	12%
31-35	18%
36-40	25%

Table 3: Effect of AGE on PEFR

Age group (in yrs.)	No. of patients (N) (300)	Mean PEFR %	No of pts. with significantly low PEFR (n= 93)
15-20	36	76	15(16.12%)
21-25	148	86	29(31.02%)
26-30	86	75	23(24.73%)

31-35	30	75	26(27.95%)
-------	----	----	------------

p<0.001,highly significant

Table 4: Effect of parity (P) on PEFR

Parity	No of cases (180)	PEFR Range	Mean PEFR (%)
G1	45	234-312	81
G2P1	45	204-278	78
G3P2	45	178-260	80
G4P3 & more	45	148-174	71

(p=0.96,NS)

Table 5: Effect of mode of termination of pregnancy (MOTP) on PEFR

Mode of termination of pregnancy	No of cases	PEFR range	PEFR Mean (%)	S/NS
Pre term delivery with episiotomy	7	168-248	76	S
Full term normal delivery with episiotomy	38	142-214	70	S
LSCS	15	126-170	66	S

Table 6: PEFR in Post natal period

PNC DAY	PEFR RANGE	PEFR MEAN (%)	S/NS
1-4	146-186	68	S
5-8	180-226	70	S
30 th day	240-356	82	NS

Discussion

The enlarged full term uterus, in later months of pregnancy, causes elevation of diaphragm by 4cm. The lower ribs flare out resulting in increase in sub costal angle and there is also increase in transverse diameter of the chest.[1,2,3]. Peak expiratory flow rate and their percentage of predicted values were significantly lower during third trimester of pregnancy compared to controls and It progressively decreased from first to third trimester. Similar results were seen in a study conducted by Memon et al , (2012).[4] The enlarged full term uterus, in later months of pregnancy, causes elevation of diaphragm (4cm). The lower ribs flare out

resulting in increase in sub costal angle and there is also increase in transverse diameter of the chest. The cardio-respiratory embarrassment is relieved during lightening , when a sense of relief of pressure symptoms is obtained due to engagement of the presenting part. It occurs 3 weeks before the expected date of delivery in primigravida. In multiparous women, the same occurs in late first stage of labour with rupture of membranes. The pulmonary changes are more marked in the presence of over distention of gravid uterus like hydramnios & multiple pregnancy. We have excluded such cases from our study to

maintain uniformity in the results. There is marginal increase in respiratory rate. The 'Tidal volume' and 'Minute ventilation' increases by 30-40%. The 'Functional Reserve Capacity' and 'Inspiratory Reserve Volume' may diminish by 20%. A study conducted by Neeraj et al (2010) states that the decrease in pulmonary function may be due to a decline in alveolar Pco₂ (caused by hyperventilation) which acts as bronchoconstrictor.[6]. There is increase predisposition of upper respiratory tract to hyperemia & congestion.[5] Alteration in blood gases is common. Po₂ increases above 100mmHg, Pco₂ decreases to 27-32mmHg, pH remains normal with increased renal bicarbonate excretion.[7] Dyspnoea & hyperventilation during pregnancy are common. Fetal well being and birth weight depend on a number of maternal and placental factors which includes maternal nutrition (height and weight), diet, rest, supplementation of Iron, calcium, proteins and the lung status of a woman. PEFR varies with age, height & sex of patient.[9,10,11]. The anaemic pregnant women showed lower PEFR when compared with PEFR of non anaemic pregnant women.[10]. Lung status of women during pregnancy is of great importance. Their efficacy should be assessed for maternal as well as fetal well being.

Conclusion

Pulmonary breathing exercises like deep inhalation and exhalation done during second trimester till delivery and postnatal period to improve patient's Spo₂ and oxygen carrying capacity of mother during labour, minimize fetal hypoxic events in Antepartum and Intrapartum period, helps the patient to bear down during labour. Well oxygenated uterus will involutes earlier.

References

1. Pandya KD, Chandwani S, Desai CA, Dadlani AG. Study of vital capacity and timed vital

- capacity in normal non-pregnant and pregnant women. J Obst Gynecol Ind, 1984; 36: 1053-1057.
2. L R Brancazio. S A Laifer & T Schartz. pregnancy & advancing gestation on peak expiratory rate. Obstetrics & gynaecology 1997;89:383. American college of obstetricians and gynecologist. Pulmonary disease in pregnancy. ACOG technical bulletin no.224.Washington DC:1996.
3. Brancazio LR, Laifer SA, Schwartz T. Peak Expiratory Flow Rate innormal pregnancy. Obstet Gynecol. 1997 ; 89 (3) : 383-386.
4. Memon et at, Change in Peak Expiratory Flow Rate in different trimesters of pregnancy ,Rawal Medical Journal, 2012; 37(3):243-246.
5. Neeraj et al, Effect Of Advanced Uncomplicated Pregnancy On Pulmonary Function Parameters Of North Indian Subjects, Indian J Physiol Pharmacol 2010 : 54(1) : 69-72.
6. Spiropoulos K, Prodromaki E, Tsapanos V. Effect of body position on PaO₂ and PaCO₂ during pregnancy. Gynecol Obstet Invest 2004 ; 58 : 22 – 25.
7. Juniper EF, Daniel EE, Roberts S, Kline PA, Hargreave FE, Newhouse MT, 1991. Effect of pregnancy on airway responsiveness and asthma severity. American Review of Respiratory Diseases 143: S78-S73.
8. Puranik BM, Kurhade GA, Kaore SB, Patwardhan SA, Kher JR. PEFR in pregnancy : an ongitudinal study. Indian J Physiol Pharmacol 1995 ; 39 (2) : 135-139.
9. Grindheim G, Toska K, Estensen ME, Rosseland LA., Changes in pulmonary function during pregnancy: a longitudinal cohort study,BJOG An International Journal of Obstetrics and Gynaecology2012 Jan;119(1):94-101.
10. Sunyal DK, Amin MR, Ahmed A, Begum S, Begum M, Rahman NJ. Peak expiratory flow rate in pregnant women. Bangladesh Soc Physiol 2007;2:20-23.

Source of support:Nil

Conflict of interest:Nil