

# Antibacterial Activity of *Justicia betonica* Linn

Swasha Khandeparker Naik<sup>1</sup>, B. L. Manjula<sup>2\*</sup>, M. Vaidagi Balaji<sup>3</sup>, Sugimani Marndi<sup>4</sup>, Sanjeet Kumar<sup>4</sup>, Rajkumari Supriya Devi<sup>4</sup>

## ABSTRACT

*Justicia betonica* L. (Acanthaceae) is widely used as a traditional folk medicinal herb. Conventionally, the plant is used to cure constipation, diarrhea, malaria, pain, stomach ache, vomiting, etc. The plant possesses many therapeutic uses such as analgesic, anti-inflammatory, antimalarial, and antimicrobial properties. The present work has been designed to document the ethnomedicinal properties, phytochemistry, and antibacterial activity of *J. betonica*. Phytochemical results showed that the presence of wide variety of biologically active compounds such as flavonoids, alkaloids, tannins, and phenolic compounds might be responsible for therapeutic values of *Justicia betonica*. Antibacterial activity showed that the inhibition was observed with the individual extracts and was effective against all bacterial strains tested. The present study highlights the importance of *J. betonica*.

**Keywords:** Acanthaceae, Antibacterial activity, Ethnomedicinal, Phytochemistry  
*Asian Pac. J. Health Sci.*, (2022); DOI: 10.21276/apjhs.2022.9.4S.16

## INTRODUCTION

Acanthaceae consists of about 4300 species belonging to 346 genera and top 12 most diverse families.<sup>[1]</sup> These plants are horticulturally important and cultivated as ornamental plant.<sup>[2,3]</sup> Characteristic features of Acanthaceae are opposite phyllotaxy of leaves and usually leaves with entire margin, stems are round to quadrangular with solitary or racemose inflorescence, bisexual flowers, large and petaloid bracts, 4–5 petals and sepals, 2–4 stamens, and superior ovary with two fused ovules. The fruit is often explosively dehiscent present inside loculicidal capsule. Seeds are usually borne on hook like retinacula, or retinacula lacking, surface smooth or roughened lacking trichomes or pubescent, and sometimes with hygroscopic trichomes that expand when moistened.<sup>[4,5]</sup> *Justicia* is an important genus of Acanthaceae, with approximately 700 species, with many unresolved species which are found in pantropical and tropical regions.<sup>[6]</sup> *Justicia* species are reported to occur in tropical to warm temperate regions of America, India, Indonesia, Southeast Asia, Malaysia, Pakistan, and Africa.<sup>[7]</sup> The genus *Justicia* is characterized by plants usually herbs or under shrubs with simple lanceolate leaves and sessile or sub sessile flowers in spikes or panicles.<sup>[8]</sup> *Justicia betonica* Linn., belonging to this genus, is a diffusely branched under shrub or an erect shrub. This plant is native to tropical Asia and tropical Africa and is distributed throughout India and Sri Lanka. In Kerala, the species is common and widely distributed in Western Ghats usually found along forest margins.<sup>[9,10]</sup>

Various parts of this plant have been used as traditional ayurvedic medicine in India as well as in other countries. The aerial part of this plant is used in diarrhea, inflammation, and swelling.<sup>[11]</sup> In India, the inflorescence extract of this plant is given orally to treat vomiting and constipation and used externally to wash hairs.<sup>[12]</sup> Leaves are crushed and applied to relieve pain and swelling.<sup>[13]</sup> Leaf and flower ash internally used for the treatment of cough, diarrhea, etc.<sup>[14]</sup> Leaf decoction is used to cure vomiting and headache. In Uganda, the leaves of *J. betonica* are used against HIV/AIDS. *J. betonica* is administered to lower cholesterol and is used to treat paralysis, ear aches, headaches, bruises, diarrhea, vomiting, constipation, pain, and inflammation and malaria in India.<sup>[15]</sup> The plant possesses analgesic, antimalarial, antimicrobial, antioxidant, and anti-inflammatory properties.<sup>[9,10]</sup>

<sup>1</sup>Department of Department of Botany, Goa University, Taleigao, Goa, India

<sup>2</sup>Department of Department of Botany, Sri Jagadguru Renukacharya College of Science, Arts and commerce, Bengaluru, Karnataka, India

<sup>3</sup>Department of Department of Botany, School of Bioscience and Technology Vellore Institute of Technology, Tamil Nadu, India

<sup>4</sup>Biodiversity and Conservation Labs., Ambika Prasad Research Foundation, Cuttack, Odisha, India

**Corresponding Author:** B. L. Manjula, Department of Botany, Sri Jagadguru Renukacharya College of Science, Arts and commerce, Bengaluru, Karnataka, India. E-mail: manjulasrivats@gmail.com

**How to cite this article:** Naik SK, Manjula BL, Balaji MV, Marndi S, Kumar S, Devi RS. Antibacterial activity of *Justicia betonica* Linn. *Asian Pac. J. Health Sci.*, 2022;9(4S):87-90.

**Source of support:** Nil

**Conflicts of interest:** None.

**Received:** 02/03/2022 **Revised:** 16/04/2022 **Accepted:** 02/05/2022

Keeping the importance of *Justicia* species, the present study was designed. It may support in standardization of plant material which could give a hand in ascertaining identity and purity of crude drugs from the present study.

## METHODOLOGY

### Collection of Plant for Experimental Works

The sample was collected and kept in poly bags tagged with the botanical name and sorted out as per standard sampling procedure and passport description.<sup>[16]</sup>

### Preparation of Extracts

Soxhlet method and percolation were adopted to obtain different extracts.<sup>[17,18]</sup> The collected experimental plant materials were dried at room temperature under shade and were powdered after drying using mechanical devices. The powdered material of the experimental plant was kept in thimble and extraction was carried out using the Soxhlet apparatus. The residue was collected and left for air drying and dried crude extracts were stored in

refrigerator for further phytochemical analysis and antibacterial activities.

### Phytochemical assays

Phytochemical analysis was carried out on different extracts of the whole plant of *J. betonica* using standard procedure to identify the bioactive compounds.<sup>[17,19,20]</sup>

#### Test for Tannins

To 5 ml of plant extract, 5 drops of 10% lead acetate were added. The formation of a light yellow precipitate indicated the presence of tannins.

#### Test for Saponins

One milliliter of the extract was boiled in 10 ml of distilled water and filtered with Whatman 42 filter paper. Five milliliters of filtrate were mixed with 2 ml of normal distilled water and shaken vigorously. Occurrence of stable persistent froth indicated the presence of saponins.

#### Test for Flavonoids

To 1 ml of the extract, few drops of dilute sodium hydroxide were added. The presence of flavonoids is indicated on production of an intense yellow color in the plant extract which became colorless on addition of 2–3 drops of 50% HCL.

#### Test for Terpenoids

0.5 g of plant extract was mixed with 2 ml of chloroform and equal volume of concentrated sulfuric acid was added. Terpenoid presence is confirmed by a reddish-brown coloration of interface.

#### Test for Phenolic Compounds

Five drops of 1% ferric chloride and 1 ml of potassium ferro cyanide was added to 2 ml of plant extract. A bluish-green solution showed the presence of phenolic compounds.

#### Test for Reducing Sugar

0.5 g of plant extract was dissolved with distilled water and filtered. The filtrate was boiled with two drops of Fehling's solution A and B for 5 min. An orange-red precipitate obtained indicated the presence of reducing sugar.

#### Test for Steroids

Two milliliters of plant extract were dissolved in 5 ml chloroform, and then, 5 ml of concentrated sulfuric acid were added. The formation of two phases (upper red and lower yellow with green fluorescence) indicated the presence of steroid.

#### Test for Alkaloids

Five milliliters of plant extract were mixed with 3 ml of aqueous HCL on water bath and then filtered. One milliliter of Dragendorff's was added to the filtrate. The occurrence of

orange red precipitate indicated the presence of alkaloids in the sample extract.

### Test for Carbonyl

Two milliliters of plant extract were added with two drops of 2, 4-dinitrophenyl hydrazine solution and thoroughly shaken. Yellow crystal formation indicated the presence of carbonyl.

### Antibacterial Activity

The extracts of experimental plant were screened for antibacterial activity against two Gram-positive bacteria *Streptococcus mutans* (Microbial Type Culture Collection [MTCC] 497) and *Streptococcus pyogenes* (MTCC 1926); three Gram-negative bacteria *Vibrio cholerae* (MTCC 3906), *Shigella flexneri* (MTCC 1457). All used MTCC bacterial strains were collected from Institute of Microbial Technology, Chandigarh. Antibacterial activity was done using slight modification of standard methods of agar well diffusion assay,<sup>[21]</sup> disk diffusion method,<sup>[22]</sup> and broth dilution assay.<sup>[23]</sup>

#### Agar Well Diffusion Assay

Agar well diffusion method<sup>[21]</sup> was followed to test the antibacterial activity of extracts of experimental plant parts against five bacterial strains. Nutrient agar plates were prepared as per instructions of manufacturer. 100 µl of nutrient broth cultures of the test microbes prepared a day before were poured over the plates uniformly and a lawn culture was prepared using a sterile spreader in a laminar hood. Wells (6 mm) were made using sterile borer. Stock solutions of samples were prepared in 100% DMSO (Sigma) and two-fold serial dilutions were made in amount of 100 µl per well at concentration of 0.25 and 0.5 mg/ml. 100 µl of samples were added by sterile syringes into the wells in three above-mentioned concentrations and allowed to diffuse at room temperature for 2 h. Plates were incubated at 35 ± 2°C for 18–24 h. Kanamycin served as standard antibiotic control. Triplicates were maintained and the experiment was repeated thrice. For each replicate, the readings (diameter of zone of inhibition in cm) were taken and the mean ± SD values (diameter of zone of inhibition) were recorded.

#### Disk Diffusion Assay

Antibacterial activity using disk diffusion assay was done using the 6 mm of disk prepared from Whatman filter paper.<sup>[22]</sup> Each extract was dissolved in dimethyl sulfoxide. The sets of dilutions (10 µg/disk and 50 µg/disk) of crude extracts and standard drugs were prepared. 6 mm disks were kept in the drugs for 12 h before placing on to the agar plates. The zones of growth inhibition around the disks were measured after 18–24 h of incubation at 37°C for bacteria. The sensitivities of the microbial species to the plant extracts were determined by measuring the sizes of inhibitory zones (including the diameter of disk) on the agar surface around the disks, and values <8 mm were considered as not active against microorganisms.

#### MIC using Broth Dilution Assay

All the extracts of experimental plant were screened for their antibacterial activity (modified method<sup>[23]</sup>). Antibacterial activity was assessed by Minimum Inhibitory Concentration (MIC) by

**Table 1:** Phytochemical test of selected experimental plants (*Justicia betonica*)

Plant name	Bioactive Compounds	Solvents					
		n-hexane	Petroleum ether	Methanol	Aqueous	Acetone	Ethanol
<i>Justicia betonica</i> (Whole plant)	Tannin	-	-	+++	-	+++	+++
	Saponin	-	-	-	-	-	-
	Flavonoids	-	-	+++	++	-	-
	Terpenoids	-	-	-	-	-	-
	Phenolic compounds	-	-	+++	+++	+++	+++
	Reducing sugar	+++	-	+++	-	+++	-
	Steroid	-	-	-	-	-	-
	Alkaloids	-	-	+++	+++	+++	-
	Carbonyl	-	-	-	-	-	-

+++ : High concentration, ++ : Mild concentration, + : Low concentration

serial dilution method. Selected colonies of aforesaid bacteria were picked off to a fresh isolation plate and inoculated in corresponding tubes containing 5 ml of trypticase soy broth. The broth was incubated for  $8 \pm 1$  h at  $35 \pm 2^\circ\text{C}$  until there was a visible growth. McFarland No.5 standard and Phosphate Buffer Saline were used to adjust the turbidity to get  $10^5$  cfu/mL.

### Data Interpretation

After the incubation, the tubes showing no visible growth after 8 h until 12 h were considered to be inhibition of bacteria which represent MIC values of a respective concentration. Inoculum control showed visible growth due to no antimicrobial agents, whereas the broth control showed no growth due to absence of bacteria. Triplicates were maintained and the experiments were repeated thrice for each replicate. The readings were taken as foresaid.

## RESULTS

The present study is based on both field and laboratory works. The crude powder of the collected experimental plant (*J. betonica*) was extracted using Soxhlet method. The extracts were used in the qualitative analysis of secondary metabolites, antimicrobial activity using agar well diffusion, disk diffusion, and MIC assay.

### Phytochemical Screening

An attempt has been made to evaluate the presence of bioactive compounds in the experimental plant through qualitative tests. The different extracts of *J. betonica* were analyzed to study the presence of secondary metabolites. The phytochemical screening revealed the presence of diverse secondary metabolites which are of highly pharmacological significance. The common secondary metabolites are present in the extracts are tannins, alkaloids, flavonoids, and phenolic compounds [Table 1].

### Evaluation of Antimicrobial Activity

#### Agar Well Diffusion

Antimicrobial activity of different extracts of the experimental plant (*J. betonica*) was determined against two Gram-positive bacteria *S. mutans* (MTCC 497) and *S. pyogenes* (MTCC 1926); three Gram-negative bacteria *V. cholerae* (MTCC 3906) and *S. flexneri* (MTCC 1457). The results revealed that the methanolic extract showed highest zone of inhibition against all the pathogens [Table 2].

**Table 2:** Antibacterial activity of *Justicia betonica* using Agar Well Diffusion method

Extracts	Zone of Inhibition (mm)			
	<i>Streptococcus mutans</i>	<i>Streptococcus pyogenes</i>	<i>Vibrio cholerae</i>	<i>Shigella flexneri</i>
Aqueous	8.5	8.0	10.6	10.5
Methanol	10.8	10.3	12.1	11.5
Acetone	9.5	9.2	10.5	10.8
Ethanol	8.2	8.1	9.5	9.3
n-hexane	8.0	8.0	9.2	9.4

**Table 3:** Antibacterial activity of *Justicia betonica* using disk diffusion method

Extracts	Zone of Inhibition (mm)			
	<i>Streptococcus mutans</i>	<i>Streptococcus pyogenes</i>	<i>Vibrio cholerae</i>	<i>Shigella flexneri</i>
Aqueous	8.5	10.5	9.5	11.5
Methanol	9.3	10.2	11.0	9.5
Acetone	8.2	8.3	9.2	9.0
Ethanol	7.2	7.5	8.1	8.0
n-hexane	7.0	7.2	8.5	8.0

#### Disk Diffusion Assay

Disk diffusion assay was carried out and it was examined that the zone of inhibition was aqueous, methanol and acetone extract for all the pathogens. It was observed that highest zone of inhibition in Aqueous extract was 11.5 mm against *S. flexneri*. Methanol, acetone, ethanol, and n-hexane extracts showed zone of inhibition of 11.0 mm, 9.2 mm, 8.1 mm, and 8.5 mm, respectively, against *V. cholerae* [Table 3].

#### Broth Dilution Assay


Broth dilution was carried out using different extracts to determine the MIC values of all the extracts against *S. mutans*, *S. pyogenes*, *Vibrio cholerae*, and *S. flexneri*. The result of MIC in aqueous, methanol, acetone, and ethanol extract was 400 mg/mm and 350 mg/mm against *V. cholerae* and *S. flexneri* [Table 4].

## DISCUSSION

*J. betonica* plays a significant role in folk as well as in veterinary medicines, but it is a less explored plant species as compared to its traditional usefulness as well as its antibacterial activity. Sasikumar et al., in 2007, reported that the extract of *J. betonica* failed to inhibit the growth of *K. pneumonia*, *S. typhi*, and *V. cholerae*. The leaf extract of this plant is reported to be active against rice moth, Coraryra.<sup>[24]</sup>

**Table 4:** Broth dilution assay of *Justicia betonica*

Extracts	MIC (mg/ml)			
	<i>Streptococcus mutans</i>	<i>Streptococcus pyogenes</i>	<i>Vibrio cholerae</i>	<i>Shigella flexneri</i>
Aqueous	500	500	400	400
Methanol	400	400	350	350
Acetone	400	400	350	350
Ethanol	450	450	400	400
n-hexane	Growth	Growth	Growth	Growth
Inoculum	Growth	Growth	Growth	Growth
Broth	No growth	No growth	No growth	No growth

**Figure 1:** Plant extracts have great potential as antimicrobial compounds against microorganisms

Ssenyondo *et al.*, in 2020, reported that the leaf extract of *J. betonica* shows antimicrobial activity against *S. aureus*.<sup>[25]</sup> According to Correa and Alcantara 2012, lignans are the major components of the active extracts of the species of *Justicia*, exhibiting important pharmacological properties, such as antiviral, antitumoral, anti-inflammatory, and antiplatelet aggregation activities, which warrant further exploration.<sup>[8]</sup> Here, we found the whole plant extract inhibits the growth of *S. mutans*, *S. pyogenes*, *V. cholerae* and *S. flexneri*.

## CONCLUSION

Since it is an important herbal remedy in traditional medicine, *J. betonica* has been extensively investigated by various researchers for its phytochemical, pharmacological, and microbiological attributes. The plant extracts have great potential as antimicrobial compounds against microorganisms [Figure 1]. Thus, they can be used in the treatment of infectious diseases caused by resistant microbes. The present study concluded that the selected medicinal plant has good activity against the clinically isolated pathogenic bacterial strains; further study is needed on the selected medicinal plant.

## REFERENCES

- Wasshausen DC, Wood JR. Contributions from the United States National Herbarium Acanthaceae of Bolivia. Vol. 49; 2004. p. 1-52.
- Amirul AA, Noraini T, Nurul AC, Ruzi AR. Petal Anatomy of Four *Justicia* (Acanthaceae) Species. Vol. 1571. AIP Conference Proceedings; 2013. p. 368-71.

- Awan A, Ahmed CB, Uzair M, Aslam MS, Farooq U, Ishfaq K. Family acanthaceae and genus *Aphelandra*: Ethnopharmacological and phytochemical review. Int J Pharm Sci 2014;6:44-55.
- Khan I, Ahmad JS, Shinwari ZK, Muhammad A, Khan Y, Kumar T. Ethnobotany and medicinal uses of folklore medicinal plants belonging to family acanthaceae: An updated review. MOJ Biol Med 2017;1:34-8.
- Manokari M, Latha R, Priyadarshini S, Cokul RM, Beniwal P, Manjunatha RY, *et al.* A comprehensive review on a less explored medicinally important plant *Justicia betonica* L. World Sci News 2019;131:110-22.
- Daniel FT. *Justicia* (acanthaceae) in texas. J Bot Res Inst Texas 2011;5:595-618.
- Perveen A, Kaiser M. Pollen flora of Pakistan-LXVII: Acanthaceae. Pak J Bot 2012;42:175-91.
- Correa GM, De Alcantara. Chemical constituents and biological activities of species of *Justicia*: A review. Rev Bras Farmacogn 2012;22:220-38.
- Eapen J, Ansary PY, Hameed AS, Indulekha VC, Resny AR. Pharmacognosy of leaf, stem and root of *Justicia betonica* Linn. Int J Res Ayurveda Pharm 2019;9:44-8.
- Hooker JD. Flora of British India. Vol. 4. United Kingdom: Reeve and Co. Ltd., Authority of the Secretary of State for India in Council; 1992. p. 525.
- Gangabhavani K, Ravishankar K. Evaluation of analgesic and anti-inflammatory activities of ethanolic extract of whole plant *Justicia betonica*. World J Pharm Pharm Sci 2013;2:5218-28.
- Rao DM, Rao UV, Sudharshanam G. Ethno-medico-botanical studies from Rayalaseema region of South Eastern Ghats, Andhra Pradesh, India. Ethnobot Leaf 2006;10:198-207.
- Khare CP. Indian Medicinal Plants: An Illustrated Dictionary. Berlin: Springer Verlag; 2007. p. 250.
- Jeruto P, Lukhoba C, Ouma G, Otieno D, Mutai C. Herbal treatments in Aldai and Kaptumo divisions in Nandi district, Rift valley Province, Kenya. Afr J Tradit Complement Altern Med 2008;5:103-5.
- Lamorde M, Tabuti JR, Obua C, Byobona CK, Lanyero H, Kibwika PB, *et al.* Medicinal plants used by traditional medicine practitioners for the treatment of HIV/AIDS and related conditions in Uganda. J Ethnopharmacol 2010;130:43-53.
- Swain J, Kumar S, Parida S, Jena PK. *In vitro* screening of antibacterial activities of selected wild tuberous plants of Odisha, India. Asian J Biol Life Sci 2020;9:79-87.
- Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and extraction: A review. Int Pharm Sci 2011;1:98-106.
- Kumar S, Mahanti P, Rath SK, Patra JK. Qualitative phytochemical analysis and antibacterial activity of *Dioscorea alata* L.: A nutraceutical tuber crops of rural Odisha. J Altern Med Res 2017;3:122-2.
- Harbone JB. Phytochemicals Method. London: Chapman and Hall Ltd.; 1973. p. 49-188.
- Trease GE, Evans WC. Pharmacognosy. London: WB Scanders Company, Ltd.; 1989. p. 89-300.
- Kumar S, Behera SP, Jena PK. Validation of Tribal claim on *Dioscorea pentaphylla* L. through phytochemical screening and evaluation of antibacterial activity. Plant Sci Res 2013;35:55-61.
- Kumar S, Jena PK. Chromatographic, antibacterial and FT-IR analysis of *Dioscorea pentaphylla* L. tuber extracts. Plant Sci Res 2014;36:83-90.
- Rai US, Arun M, Shetty I, Vijesh AM, Prabhu N, Isloor S. Novel chromeno (2,3-b) pyrimidine derivatives as potential anti-microbial agents. Eur J Med Chem 2010;45:2695-9.
- Sasikumar JM, Thayumanavan T, Subashkumar R, Janardhanan K, Lakshmanaperumalsamy P. Antibacterial activity of some ethnomedicinal plants from the Nilgiris, Tamil Nadu, India. Nat Prod Radiance 2007;6:34-9.
- Ssenyondo KA, Tibyangye J, Kinene MA, Christopher E, Stephen K, Winnie N, *et al.* Antimicrobial activity of *Vernonia amygdalina*, *Justicia betonica*, *Leonatis nepetaefolium* and *Momordica foetida* to *Staphylococcus aureus*. J Pharm Biol Sci 2010;15:6-10.