

Synthesis, Characterization and Antimicrobial evaluation of novel Azole Based (p-Toluic Acid) derivatives**Omprakash Sharma^{1*}, Pankaj Sharma², Birendra Shrivastava³, Jitender Singh⁴**¹Research Scholar, School of Pharmaceutical Sciences, Jaipur National University, Jaipur, Rajasthan, India²Professor, School of Pharmaceutical Sciences, Jaipur National University, Jaipur, Rajasthan, India³Director, School of Pharmaceutical Sciences, Jaipur National University, Jaipur, Rajasthan, India⁴Professor, Lord Shiva College of Pharmacy, Sirsa, Haryana, India

Received: 15-06-2018 / Revised: 10-09-2018 / Accepted: 25-09-2018

Abstract

This study includes the Synthesis of Azoles Based p-Toluic Acid Derivative. The compounds were prepared by reaction of p-Toluic Acid with thiosemicarbazide in the presence of Phosphorus oxychloride to yield 1, 3, 4-thiadiazole nucleus. Further this thiadiazole was treated with different aromatic aldehydes in presence of methanol to yield corresponding Schiff's bases. All the Schiff's bases thus obtained were reacted with Thioglycolic acid to yield the title compounds. Synthesized derivatives were characterized by IR, ¹H NMR and screened for their antibacterial (*S. aureus*, *B. subtilis*, *E. coli*, and *P. aeruginosa*) and antifungal activity (*A. Niger* and *C. albicans*).

Key Words: Azole derivative, p-Toluic Acid derivative, antimicrobial activity**Introduction**

Medicine is a system of scientific knowledge and practical activities aimed at preserving and improving the health of people, prevention and treatment of human disease. [1]. This would serve as the basis for ensuring good health for a community Traditional Indian medicine-a conglomerate of Ayurveda, Sidha and Unani has a long history The 4-thiazolidinones that do not contain aryl or higher alkyl substituents are somewhat soluble in water or thiazolidinone that do not contain aryl or higher alkyl substituents are somewhat less soluble in water[2] Thiadiazole is a heterocyclic compound featuring both a nitrogen and sulphur atom as a part of the aromatic five membered rings. It acts as Hydrogen bonding domain and two electron donor moiety along with ability of Bio-isosteric replacement of thiazole moiety. Bioisosteres are atoms or group of molecules that fit the broadest definition for isosteres. [3-5].

Thiazole is resistant to oxidizing agents; even hot nitric acid has little effect. It is either unaffected by reducing agents or at most some ring opening occurs it is therefore not possible to make dihydrothiazoles or thiazolidines by catalytic dehydrogenation but they must need to be prepared by means of condensation reactions. Thiazole is of great biological importance as it is used as an intermediate to manufacture synthetic drugs, fungicides and dyes. A thiazole ring is found in the essential vitamin B₁ [7]. The literature survey showed that the Thiadiazole moiety exhibit antimicrobial, anti cancer, anti inflammatory, antioxidant Anti Fungal, anti depressant, anti-Anthelmintic activity. [8-13]. Considerable confusion concerning the structure of 4-thiazolidinones exist in the early literature and non cyclic formulas were at first proposed for pseudo thiohydantoin and for rhodanine 1,4-thiazolidinones are derivatives of thiazolidine with a carbonyl group at the 4th position. Substitution is possible at 2, 3 and 5 positions various optical and geometrical isomers are reported [19]. The initial synthetic route is not economical enough for providing sufficiently quantity of the 1,3-thiazolidin-4-one intermediate and the authors developed an improvement by protection of alanine with chloro trimethylsilane (TMSCl). The direct condensation of aldehydes, alanine and mercaptoacetic acid give the

Correspondence*Omprakash Sharma**

Research Scholar, School of Pharmaceutical Sciences, Jaipur National University, Jaipur, Rajasthan, India and Lecturer, Government Polytechnic, Mandi Adampur, Hisar, India.

Mobile No- +91-9466746250, 9466869515

E-Mail: sharmaop1873@gmail.com


desired thiazolidinone along with the dithioacetal as the major byproduct [20].

Materials and Methods

The chemical used for experimental were of synthetic grade and procured from Nice Laboratory, Loba Chemie Private Limited, CDH New Delhi and Research Lab Fine Chem industries Mumbai. The melting points of the synthesized compounds were determined in open glass capillaries and are uncorrected. IR spectra were recorded on FTIR

Spectrometer. ^1H NMR spectra were recorded on were recorded in $\text{CDCl}_3/\text{DMSO}$ solution on a Bruker Avance II 400 MHz NMR spectrometer taking Tetramethylsilane (TMS) as internal standard. Progress of reaction was monitored by TLC using solvent Hexane: Ethyl acetate (40%) & Methanol: Chloroform (1%).

Scheme



Ar =(a)= 4-Cl- C_6H_5^- , (b) 2-F- C_6H_5^- , (c) 2- NO_2 - C_6H_5^- , (d) 3-OH- C_6H_5^- , (e) 4-OH- C_6H_5^- , (f) 2-O CH_3 - C_6H_5^- ,
(g) 4-O CH_3 - C_6H_5^- (h) 4-OH- C_6H_5^- (i) 2,4-Cl- C_6H_5^- (j) 4- NO_2 - C_6H_5^- (k) 2-Br- C_6H_5^-

Synthetic procedure

Synthesis of 5-p-tolyl-1, 3, 4-thiadiazol-2-amine

A mixture of equimolar quantity of p-Toluic Acid (50mmol) and thiosemicarbazide (50 mmol) were refluxed in 20 ml POCl_3 solution for 2 h at 75°C . After cooling down to room temperature, cold water was added. The mixture was again refluxed for 4 h. The content was cooled down at room temperature and made alkaline at pH 8.0 by addition of 50 % NaOH solution. The solid (5-p-tolyl-1, 3, 4-thiadiazol-2-amine) (compound 3) so obtained was filtered and recrystallized using ethanol. (Bhatia et al)

Synthesis of Schiff bases

In this step compound 3 i.e. 5-p-tolyl-1, 3, 4-thiadiazol-2-amine was refluxed with different substituted aromatic aldehydes (0.025M) [a-k] using methanol (50 mL) as solvent in presence of small amount of glacial acetic acid for 2 hrs. The mixture was cooled and poured in ice water. The content was put undisturbed for 24h. The solid thus obtained was separated by filtration and recrystallized from Methanol to give the Corresponding Schiff bases (a-k). [Kumar et al 2011 and Kumar et al 2018]

Synthesis of thiazolidinone

A mixture of Synthesized Schiff base (0.02M) (a-k) and Thioglycolic acid (0.02M) in 50ml DMF,

containing a pinch of anhydrous $ZnCl_2$ were refluxed for about 6 hrs. The reaction mixture was cooled and poured on to crushed ice. The solid thus obtained was filtered, washed with water and the product thiazolidinone (6a-k) was recrystallized from ethanol. [Sharma et al 2010].

Characterization of synthesized compounds

Compound 1

2-(4-chloro-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3014.1 (C-H str, aromatic), 2985.5 (C-H str, alkane), 1674.4 (C=O str.), 1636.9 (C=N str.), 1284.1 (C-N), 835.40(C-Cl).

1H NMR (DMSO) (δ -ppm): 7.05-7.55 (8H, Ar H); 5.7 (1H, N-CH); 3.6 (1H, CH-S); 2.31(3H, s, CH_3).

Compound 2

2-(2-Fluoro-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3012.8 (C-H str, aromatic), 2987.4 (C-H str, alkane), 1671.6 (C=O str.), 1637.5 (C=N str.), 1284.6 (C-N)

1H NMR (DMSO) (δ -ppm): 6.9-7.45 (8H, Ar H); 5.74 (1H, N-CH); 3.45 (1H, CH-S); 2.28(3H, s, CH_3)

Compound 3

2-(2-Nitrophenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3007.1 (C-H str, aromatic), 2988.4 (C-H str, alkane), 1670.6 (C=O str.), 1636.3 (C=N str.), 1285.0 (C-N)

1H NMR (DMSO) (δ -ppm): 7.2-7.85 (8H, Ar H); 5.9 (1H, N-CH); 3.47 (1H, CH-S); 2.30(3H, s, CH_3)

Compound 4

2-(3-hydroxy-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}):3385.18 (O-H str.), 3009.05 (C-H str, aromatic), 2981.56 (C-H str, alkane), 1668.62 (C=O str.), 1635.54 (C=N str.), 1280.74 (C-N)

1H NMR (DMSO) (δ -ppm): 6.78-7.51(8H, m, Ar-H); 5.88 (1H, s, N-CH); 4.8 (1H, s, OH), 3.44 (1H,s, CH-S); 2.26 (3H, s, CH_3)

Compound 5

2-(4-Hydroxy-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}):3384.42 (O-H str.), 3010.12 (C-H str, aromatic), 2981.09 (C-H str, alkane), 1667.84 (C=O str.), 1635.31 (C=N str.), 1281.04 (C-N)

1H NMR (DMSO) (δ -ppm): 6.74-7.48 (8H,m, Ar-H); 5.86 (1H, s, N-CH); 4.91 (1H,s, OH), 3.42 (1H,s, CH-S); 2.27 (3H, s, CH_3)

Compound 6

2-(2-Methoxy-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3018.10 (C-H str, aromatic), 2978.29 (C-H str, alkane), 1668.08 (C=O str.), 1637.49 (C=N str.), 1285.04 (C-N)

1H NMR (DMSO) (δ -ppm): 6.87-7.62 (8H, m, Ar-H); 5.89 (1H, s, N-CH); 3.76 (3H,s,-OCH₃), 3.45(1H, s,CH-S); 2.29 (3H, s, CH_3)

Compound 7

2-(4-Methoxy-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3016.2 (C-H str, aromatic), 2978.45 (C-H str, alkane), 1669.18 (C=O str.), 1636.11 (C=N str.), 1283.47 (C-N)

1H NMR (DMSO) (δ -ppm): 6.85-7.71 (8H, m, Ar-H); 5.86 (1H, s, N-CH); 3.78 (3H,s, -OCH₃), 3.41(1H,s, CH-S); 2.33 (3H, s, CH_3)

Compound 8

2-Phenyl-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3013.8 (C-H str, aromatic), 2977.95 (C-H str, alkane), 1666.09 (C=O str.), 1631.65 (C=N str.), 1277.06 (C-N)

1H NMR (DMSO) (δ -ppm): 7.08-7.82 (9H, m, Ar-H); 5.86 (1H, s, N-CH); 3.78 (3H, s, -OCH₃), 3.41(1H, s, CH-S); 2.33(3H, s, CH_3)

Compound 9

2-(2, 4-Dichloro-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3022.03 (C-H str, aromatic), 2979.05 (C-H str, alkane), 1668.11 (C=O str.), 1630.25 (C=N str.), 1278.94 (C-N)

1H NMR (DMSO) (δ -ppm): 7.05-7.81 (7H, m, Ar-H); 5.85 (1H, s, N-CH); 3.49 (1H, s, CH-S); 2.35(3H, s, CH_3)

Compound 10

2-(4-Nitro-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3020.0 (C-H str, aromatic), 2982.24 (C-H str, alkane), 1673.52 (C=O str.), 1635.74 (C=N str.), 1277.5 (C-N)

1H NMR (DMSO) (δ -ppm): 7.15-8.01 (8H, m, Ar-H); 5.89 (1H, s, N-CH); 3.37 (1H, s, CH-S); 2.28(3H, s, CH_3)

Compound 11

2-(2-Bromo-phenyl)-3-(5-p-tolyl-[1, 3, 4]-thiadiazol-2-yl)-thiazolidin-4-one

IR (KBr) (cm^{-1}): 3021.08 (C-H str, aromatic), 2978.51 (C-H str, alkane), 1670.14 (C=O str.), 1634.56 (C=N str.), 1274.6 (C-N)

1H NMR (DMSO) (δ -ppm): 7.02-7.96 (8H, m, Ar-H); 5.83 (1H, s, N-CH); 3.39 (1H, s, CH-S); 2.29(3H, s, CH_3).

Biological Activity

Keeping in view the activity profile of Thiazolidinone and Thiadiazole itself, it was thought sensible to carry out biological screening of all the newly synthesized compounds. Therefore all the synthesized compounds were subjected to Antimicrobial evaluation.

Antibacterial activity

The antibacterial activities of compounds were determined *in vitro* against Gram-positive (*Staphylococcus aureus*, *Bacillus subtilis*) and Gram-negative (*Escherichia coli*, *P. aeruginosa*) bacteria by tube dilution method. Double strength nutrient broth (I.P.) media and sabourand's glucose broth media (I.P.) were used for bacteria and fungi respectively.

The samples were incubated at 37 °C for 24 h (bacteria), at 25 °C for 7 d (*A. Niger*) and at 37 °C for 48 h (*C. albicans*) and the results were recorded in terms of minimum inhibitory concentration (MIC). Ciprofloxacin and fluconazole were taken as standard drugs for antibacterial and antifungal activity, respectively.

Determination of MIC

MIC of compounds was determined by two fold serial dilution technique. Dilution of test compound and standard drugs were prepared in test medium to give a concentration of 50, 25, 12.5, 6.25, 3.125 and 1.56 µg/ml from stock solution (100µg/ml). All the samples were inoculated with 0.1ml suspension of bacteria in saline and incubated at required temperature. MIC was determined by lowest concentration of sample that prevents the development of turbidity. [18] The observed MIC is presented in Table 1.

Table 1: Minimum Inhibitory Concentration (MIC) of different compounds

Compounds	Minimum Inhibitory Concentration (MIC)					
	Bacterial strains				Fungal strains	
	Gram Positive		Gram Negative			
	<i>S. aureus</i>	<i>B. subtilis</i>	<i>P. aeruginosa</i>	<i>E.coli</i>	<i>C. albicans</i>	<i>A. niger</i>
1	1.610	1.610	1.610	1.610	1.610	1.610
2	0.842	0.842	0.842	0.842	0.842	0.842
3	1.570	1.570	0.785	0.785	1.570	1.570
4	1.689	1.689	1.689	1.689	1.689	1.689
5	1.689	1.689	1.689	1.689	1.689	1.689
6	1.74	1.74	1.74	1.74	0.870	0.870
7	1.74	1.74	1.74	1.74	0.870	0.870
8	1.77	1.77	1.77	1.77	1.77	1.77
9	0.740	0.740	0.740	0.740	1.48	1.48
10	1.570	1.570	0.785	0.785	1.570	1.570
11	0.723	0.723	0.723	0.723	0.723	0.723
Std.	0.471	0.471	0.471	0.471	0.510	0.510












Standard: ciprofloxacin (antibacterial) and fluconazole (antifungal).

Result and discussion

The Title compounds were synthesized as per the procedure mentioned under heading Synthetic procedure. The synthesized compounds were recrystallized and purity of the compounds was ascertained by using appropriate solvent systems (ethyl acetate: benzene: water). The compounds were characterized by their IR and NMR spectroscopic data.

The result of IR and HNMR data are in good agreement with their theoretical values. The Physiochemical data of Synthesized Thiazolidinone derivatives is presented in table 2. Compounds 2, 3, 6, 7,9,10, 11 showed the significant antimicrobial activity against different strains of microbes.

Table 2: Characterization (Physiochemical) data of compounds

Compound no	Ar-	Molecular formula	Molecular weight	Melting point	% yield
1		C ₁₈ H ₁₄ ClN ₃ OS ₂	387.91	200-205	74
2		C ₁₈ H ₁₄ FN ₃ OS ₂	371.45	210-212	69
3		C ₁₈ H ₁₄ N ₄ O ₃ S ₂	398.46	180-184	66
4		C ₁₈ H ₁₅ N ₃ O ₂ S ₂	369.46	180-184	78
5		C ₁₈ H ₁₅ N ₃ O ₂ S ₂	369.46	182-186	70
6		C ₁₉ H ₁₇ N ₃ O ₂ S ₂	383.49	170-174	75
7		C ₁₉ H ₁₇ N ₃ O ₂ S ₂	383.49	168-170	80
8		C ₁₈ H ₁₅ N ₃ OS ₂	353.46	180-182	74
9		C ₁₈ H ₁₃ Cl ₂ N ₃ OS ₂	422.35	180-184	60
10		C ₁₈ H ₁₄ N ₄ O ₃ S ₂	398.46	190-192	75
11		C ₁₈ H ₁₄ BrN ₃ OS ₂	432.36	190-193	62

Conclusion

In the present work two azole containing moiety i.e. Thiadiazole and Thiazolidinone were clubbed to obtain their synergistic effect and evaluated for their antimicrobial effect. The experimental data of the present research work reveals that synthesized azole derivatives of demonstrate significant Antibacterial and Antifungal Activity. Compound 2, 3, 6, 7,9,10, 11

showed significant antimicrobial activity (table 1) Further, it has been observed that Thiadiazole which are substituted by halogen (-Cl, -Br, -F) with electron withdrawing property exhibits more Antimicrobial activity than the other.

References

1. Patwardhan B, Vaidya A, Chorghade M. Ayurveda and natural products drug discovery. *Curr. Sci.*, 2004; 86(6):789-799.

2. Mulay, A.; Ghodke, M.; Nikalje, PA. Exploring potential of 4- thiazolidinone: A brief review. *Int J. Pharm. Pharm. Sci.*, 2009; 1(1): 47-64.
3. Ogundele S.O. Aspects of indigenous medicine in south western Nigeria. *Ethno. Med.*, 2007; 1(2): 127-133.
4. Siddiqui N, Ahuja P, W. Ahsan W, Pandeya and Alam M.S. Thiadiazole: progress report on Biological Activities. *J. chem. Pharm. Res.* 2009;1:19-30.
5. Priyanka L. Gaikwad, Priyanka S. Gandhi, Deepali M. Jagdale, Vilasrao J. Kadam. The Use of Bioisosterism in Drug Design and Molecular Modification. *Am. J. PharmTech Res.* 2012; 2(4):1-23.
6. Al-Amiery A.A.H, Yousif M, Shakir M.A. Synthesis of novel metal complexes of 5-(4-Isopropoxyphenyl)-Nphenyl-1,3,4-thiadiazole-2-amine, *Eng .Tech. J.* 2009; 27: 891-897.
7. Singh T, Sharma PK, Mondal SC, Kumar N. Synthesis, anticonvulsant activity & spectral characterization of some novel thiazolidinone derivatives. *J. Chem. Pharm. Res.*, 2011; 3(5): 609-615.
8. Patel N.B, Shaikh F.M. Synthesis and antimicrobial activity of new 4-thiazolidinone derivatives containing 2-amino-6-methoxybenzothiazole. *Saudi Pharm. J.*, 2010; 18(3): 129-136.
9. Srivastava SK, Srivastava S, Srivastava SD. Synthesis and biological significance of 2-mercaptobenzoxazole derivatives. *J. Chem. Pharm. Res.*, 2010; 2(5): 449-457.
10. Chatrabhuji PM, Nimavat KS, Vyas KB, Undavia NK. Biological studies of some novel 2-aryl-3-[(4-methyl cinnamoyl amino)-4-oxo-thiazolidines. *Der Pharma Chemica.*, 2010; 2(3): 109-113.
11. Bhatt JJ, Shah BR, Shah HP, Trivedi PB. Synthesis, anticonvulsant activity and spectral characterization of some novel thiazolidinone derivatives. *J. Chem. Pharm. Res.*, 1994; 25: 189-192.
12. Shih MH, Fy K. Synthesis and evaluation of antioxidant activity of substituted thiazolidinone and thiazoline derivatives. *Bioorg & Med Chem.*, 2004; 12: 4633-4644.
13. Havrylyuk, D, Zimenkovsky B, Vasylenko O, Zaprutko L, Gzella A. Synthesis of novel thiazolone-based compounds containing pyrazoline moiety and evaluation of their anticancer activity. *Eur. J. Med. Chem.*, 2009; 44(4): 1396-1404.
14. Bhatia Rohit, Kaur Amandeep. Synthesis spectral studies and antimicrobial activity of some imidazole [2,1-b][1,3,4] thiadiazole derivatives. *Der Pharma Chemica*, 2014; 6(6):114-120.
15. Kumar V, Sharma A. and Sharma P.C. Synthesis of some novel 2,5-disubstituted thiazolidinones from a long chain fatty acid as possible antiinflammatory, analgesic and hydrogen peroxide scavenging agents. *J. Enz. Inhib. Med. Chem.*, 2011; 26(2): 198-203.
16. Sharma A, Kumar V, Sharma PC. J. Synthesis and Biological evaluation of novel benzothiazole clubbed fluoroquinolone derivatives. *Enz. Inhib. Med. Chem., J. Enz. Inhib. Med. Chem.*, 2011; 26(4): 546-552.
17. Aakashdeep, Jain S, Sharma P.C. Synthesis and anti-inflammatory activity of some novel biphenyl-4-carboxylic acid 5-(arylidene)-2-(aryl)-4-oxothiazolidin-3-yl amides. *Acta Pol. Pharm. Drug Res.*, 2010; 67(1): 63-67.
18. Kumar Ramesh, Kumar Mahesh. Synthesis of 3-[4-(2-Amino-6 (substituted phenyl)-pyrimidin-4-yl)-phenylimino]-5-chloro-1, 3-dihydro-indol-2-one derivatives of 5-chloroisatin as potential antimicrobial agents. *J Pharm Chem Biol Sci* , 2018; 5(4):399-404.
19. Drews, J. *Drug Discovery: A historical perspective.* Drug Design, 1st Ed., Nirali Prakashan, Pune, 1995; 4(1): 1-15.
20. Kato, T.; Ozaki, T.; Suzuki, Y.; Ohi, N. Novel calcium antagonists with both calcium overload inhibition and antioxidant activity. *J. Med. Chem.*, 1998; 41(2): 4309-16.

Conflict of Interest: None

Source of Support: Nil